


## AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions and listings of claims in the application:

**Listing of Claims:**

1. (Currently Amended) A compound of formula I



wherein

X is =CR<sup>0</sup>- or =N-;

each of R<sup>0</sup>, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> independently is hydrogen; hydroxy; C<sub>1</sub>-C<sub>8</sub>alkyl; C<sub>2</sub>-C<sub>8</sub>alkenyl; C<sub>3</sub>-C<sub>8</sub>cycloalkyl; C<sub>3</sub>-C<sub>8</sub>cycloalkyl-C<sub>1</sub>-C<sub>8</sub>alkyl; hydroxyC<sub>1</sub>-C<sub>8</sub>alkyl; C<sub>1</sub>-C<sub>8</sub>alkoxyC<sub>1</sub>-C<sub>8</sub>alkyl; hydroxyC<sub>1</sub>-C<sub>8</sub>alkoxyC<sub>1</sub>-C<sub>8</sub>alkyl; arylC<sub>1</sub>-C<sub>8</sub>alkyl which optionally may be substituted on the ring by hydroxy, C<sub>1</sub>-C<sub>8</sub>alkoxy, carboxy or C<sub>1</sub>-C<sub>8</sub>alkoxycarbonyl;

~~or R<sup>3</sup> and R<sup>4</sup> form together with the nitrogen and carbon atoms to which they are attached a 5 to 10 membered heterocyclic ring and having 1, 2 or 3 heteroatoms selected from N, O and S;~~

or each of R<sup>1</sup>, and R<sup>2</sup> ~~and R<sup>3</sup>~~, independently, is halogen; halo-C<sub>1</sub>-C<sub>8</sub>alkyl; C<sub>1</sub>-C<sub>8</sub>alkoxy; halo-C<sub>1</sub>-C<sub>8</sub>alkoxy; hydroxyC<sub>1</sub>-C<sub>8</sub>alkoxy; C<sub>1</sub>-C<sub>8</sub>alkoxyC<sub>1</sub>-C<sub>8</sub>alkoxy; aryl; arylC<sub>1</sub>-C<sub>8</sub>alkoxy; heteroaryl; heteroaryl-C<sub>1</sub>-C<sub>4</sub>alkyl; 5 to 10 membered heterocyclic ring; nitro; carboxy;

C<sub>2</sub>-C<sub>8</sub>alkoxycarbonyl; C<sub>2</sub>-C<sub>8</sub>alkylcarbonyl; -N(C<sub>1</sub>-C<sub>8</sub>alkyl)C(O) C<sub>1</sub>-C<sub>8</sub>alkyl; -N(R<sup>10</sup>)R<sup>11</sup>;

-CON(R<sup>10</sup>)R<sup>11</sup>; -SO<sub>2</sub>N(R<sup>10</sup>)R<sup>11</sup>; or -C<sub>1</sub>-C<sub>4</sub>-alkylene-SO<sub>2</sub>N(R<sup>10</sup>)R<sup>11</sup>; wherein each of R<sup>10</sup> and R<sup>11</sup> independently is hydrogen; hydroxy; C<sub>1</sub>-C<sub>8</sub>alkyl; C<sub>2</sub>-C<sub>8</sub>alkenyl; C<sub>3</sub>-C<sub>8</sub>cycloalkyl;

C<sub>2</sub>-C<sub>8</sub>cycloalkyl-C<sub>1</sub>-C<sub>8</sub>alkyl; C<sub>1</sub>-C<sub>8</sub>alkoxyC<sub>1</sub>-C<sub>8</sub>alkyl; hydroxyC<sub>1</sub>-C<sub>8</sub>alkoxyC<sub>1</sub>-C<sub>8</sub>alkyl; hydroxyC<sub>1</sub>-C<sub>8</sub>alkyl; (C<sub>1</sub>-C<sub>8</sub>alkyl)-carbonyl; arylC<sub>1</sub>-C<sub>8</sub>alkyl which

- optionally may be substituted on the ring by hydroxy, C<sub>1</sub>-C<sub>8</sub>alkoxy, carboxy or C<sub>2</sub>-C<sub>8</sub>alkoxycarbonyl; or 5 to 10 membered heterocyclic ring;
- or R<sup>1</sup> and R<sup>2</sup> form together with the C-atoms to which they are attached aryl or a 5 to 10 membered heteroaryl group having one or two heteroatoms selected from N, O and S; or
- R<sup>5</sup> is hydrogen; halogen; cyano; C<sub>1</sub>-C<sub>8</sub>alkyl; halo-C<sub>1</sub>-C<sub>8</sub>alkyl;
- C<sub>2</sub>-C<sub>8</sub>alkenyl; C<sub>2</sub>-C<sub>8</sub>alkynyl; C<sub>3</sub>-C<sub>8</sub>cycloalkyl; C<sub>3</sub>-C<sub>8</sub>cycloalkylC<sub>1</sub>-C<sub>8</sub>alkyl; C<sub>5</sub>-C<sub>10</sub>arylC<sub>1</sub>-C<sub>8</sub>alkyl;
- R<sup>6</sup> is hydrogen;
- each of R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> is independently hydrogen; hydroxy; C<sub>1</sub>-C<sub>8</sub>alkyl; C<sub>2</sub>-C<sub>8</sub>alkenyl; halo-C<sub>1</sub>-C<sub>8</sub>alkyl; C<sub>1</sub>-C<sub>8</sub>alkoxy; C<sub>3</sub>-C<sub>8</sub>cycloalkyl; C<sub>3</sub>-C<sub>8</sub>cycloalkylC<sub>1</sub>-C<sub>8</sub>alkyl; arylC<sub>1</sub>-C<sub>8</sub>alkyl; -Y-R<sup>12</sup> wherein Y is a direct bond or O and R<sup>12</sup> is a substituted or unsubstituted 5, 6 or 7 membered heterocyclic ring comprising 1, 2 or 3 heteroatoms selected from N, O and S; carboxy; (C<sub>1</sub>-C<sub>8</sub>alkoxy)-carbonyl; -N(C<sub>1</sub>-C<sub>8</sub>alkyl)-CO-NR<sup>10</sup>R<sup>11</sup>; -CONR<sup>10</sup>R<sup>11</sup>; -N(R<sup>10</sup>)(R<sup>11</sup>); or R<sup>7</sup> and R<sup>8</sup> or R<sup>8</sup> and R<sup>9</sup>, respectively form together with the carbon atoms to which they are attached, a 5 or 6 membered heteroaryl comprising 1, 2 or 3 heteroatoms selected from N, O and S; or a 5 or 6 membered carbocyclic ring;
- ~~provided that one of R<sup>1</sup>, R<sup>3</sup> or R<sup>3</sup> is -CON(R<sup>10</sup>)R<sup>11</sup> or -SO<sub>2</sub>N(R<sup>10</sup>)R<sup>11</sup>;~~
- in free form or salt form;
- wherein
- aryl represents phenyl, naphthyl or 1,2,3,4-tetrahydronaphthyl,
- heteroaryl is a 5 or 6 membered aromatic heterocyclic ring, optionally condensed to 1 or 2 benzene rings and/or to a further heterocyclic ring, and
- wherein a heterocyclic ring is a 5 or 6 membered heterocyclic ring being saturated or unsaturated and optionally condensed to 1 or 2 benzene rings and/or to a further heterocyclic ring
2. (Original) A process for the production of a compound of formula I according to claim 1, comprising the steps of reacting a compound of formula II wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup> and X are as defined in claim 1, and Y is a leaving group; with a compound of formula III wherein R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined in

claim 1; and recovering the resulting compound of formula I in free form or in salt form, and, where required, converting the compound of formula I obtained in free form into the desired salt form, or vice versa.

3. (Cancelled).
4. (Original) A pharmaceutical composition comprising a compound of formula I according to claim 1 or a pharmaceutically acceptable salt thereof, together with one or more pharmaceutically acceptable carriers or diluents therefor.
- 5-7. (Cancelled).
8. (Previously Presented) A method for treating breast cancer, comprising:  
administering to a subject in need thereof, a therapeutically effective amount of a compound of formula I according to claim 1 or a pharmaceutically acceptable salt thereof.
9. (Cancelled).